

# Saturation and hadronic cross sections at very high energies\*

F. CARVALHO, F.O. DURÃES, F.S. NAVARRA

Instituto de Física, Universidade de São Paulo  
C.P. 66318, 05315-97, São Paulo, SP, Brazil

AND

V.P. GONÇALVES

Instituto de Física e Matemática,  
Universidade Federal de Pelotas, Caixa Postal 354, CEP 96010-900, Pelotas, RS,  
Brazil

We propose a simple model for the total  $pp/p\bar{p}$  cross section, which is a generalization of the minijet model with the inclusion of a window in the  $p_T$ -spectrum associated to the saturation physics. Our model implies a natural cutoff for the perturbative calculations which modifies the energy behavior of this component, so that it satisfies the Froissart bound. Including the saturated component, we obtain a satisfactory description of the very high energy experimental data.

PACS numbers: 12.38.Lg, 13.60.Hb, 13.85.-t, 13.85.Dz

Long ago a QCD based explanation for the growth of the hadronic cross sections was proposed by Gaisser and Halzen [1]. In their approach, called minijet model, the total cross section can be decomposed as  $\sigma_{tot} = \sigma_0 + \sigma_{pQCD}$  where  $\sigma_0$  characterizes the nonperturbative contribution and  $\sigma_{pQCD}$  is calculable in perturbative QCD. Unfortunately, this approach implies a power-like energy behavior for the total cross section, violating the Froissart bound. Several attempts were made to reduce this too fast growth [2].

At high energies the small- $x$  gluons in a hadron wavefunction should form a Color Glass Condensate (CGC) [3]. This new state of matter is characterized by gluon saturation and by a typical momentum scale, the saturation scale  $Q_s$ , which determines the critical line separating the linear and saturation regimes of the QCD dynamics.

---

\* Presented at the “School on QCD, Low x physics, saturation and diffraction”, Copanello (Calabria, Italy), July 1 - 14 2007.




Fig. 1. Perturbative (solid lines) and saturated components (dashed lines) of the total cross section (normalized by  $\ln^2 s$ )

Some attempts to reconcile the QCD parton picture with the Froissart limit using saturation physics were proposed in recent years [4]. Here we generalize the minijet model assuming the existence of a saturation window between the nonperturbative and perturbative regimes of QCD, which grows when the energy increases, since  $Q_s$  grows with the energy. The cross section is then written as:

$$\sigma_{tot} = \sigma_0 + \sigma_{sat} + \sigma_{pQCD} \quad (1)$$

where the saturated component,  $\sigma_{sat}$ , contains the dynamics of the interactions at scales lower than the saturation scale. In our approach *the saturation scale is a cutoff at low transverse momenta of the perturbative cross section*,  $\sigma_{pQCD}$ , which is given by:

$$\sigma_{pQCD} = \frac{1}{2} \int_{Q_s^2} dp_T^2 \sum_{i,j} \int dx_1 dx_2 f_i(x_1, p_T^2) f_j(x_2, p_T^2) \hat{\sigma}_{ij} \quad (2)$$

where  $f_i(x, Q^2)$  is the parton density of the species  $i$ , with fractional momentum  $x_1$  (or  $x_2$ ) in the proton and  $\hat{\sigma}_{ij}$  is the elementary parton-parton cross section. We have used the MRST parton distributions [5]. The saturation scale is given by  $Q_s^2(x) = Q_0^2 (\frac{x_0}{x})^\lambda$ , where the parameters  $Q_0^2 = 0.3 \text{ GeV}^2$  and  $x_0 = 0.3 \times 10^{-4}$  were fixed by fitting the  $ep$  HERA data. Following [6] we take  $x = \frac{q_0^2}{s}$  and  $q_0 = 1.4 \text{ GeV}$ . Therefore we have

$$Q_s^2(s) \propto s^\lambda$$

In Fig. 1 we show in arbitrary units the energy behavior of the ratio  $\sigma_{pQCD}/\ln^2 s$  (solid lines) and  $\sigma_{sat}/\ln^2 s$  (dashed lines) for two choices of  $\lambda$ . As it can be seen the choice  $\lambda = 0.25$  leads to a fast growth of  $\sigma_{pQCD}$  until  $\sqrt{s} = 10^4$  GeV. From this point on, it grows slower than  $\ln^2 s$ . A slight increase in  $\lambda$  ( $= 0.3$ ) is enough to tame the growth of  $\sigma_{pQCD}$  already at  $\sqrt{s} \simeq 10^3$  GeV. On the other hand, a decrease in  $\lambda$  ( $= 0.1$ ) would postpone the fall of the ratio to very high energies  $\sqrt{s} \simeq 10^6$  GeV. Although the energy at which the behavior of the cross section becomes “sub-Froissart” may depend on  $\lambda$ , one conclusion seems very robust: *once  $\lambda$  is finite, at some energy the growth of the cross section will become weaker than  $\ln^2 s$ .*

For the saturated component we shall use the model proposed in Ref. [6]:

$$\sigma_{sat} = \int d^2 r_{\perp} |\Psi_p(r_{\perp})|^2 \sigma_{dip}(x, r_{\perp}) \quad (3)$$

where the proton wave function  $\Psi_p$  is chosen to be a gaussian with the typical size of the proton [7] and the dipole-proton cross section reads:

$$\sigma_{dip}(r_{\perp}, x) = 2 \int d^2 b \mathcal{N}(x, r_{\perp}, b) . \quad (4)$$

We take the dipole scattering amplitude from [8] (we call it IIM) and, following [6], introduce the  $b$  dependence by witting:

$$\mathcal{N}(x, r_{\perp}, b) = 1 - e^{-\kappa \frac{S(b)}{S(0)}} \quad (5)$$

where the parameter  $\kappa$  is related to the  $b = 0$  solution through  $\kappa = -\ln[1 - \mathcal{N}(b = 0)]$ . In (5), the profile function is assumed to be  $S(b) = e^{(-b^2/R_p^2)}$ , where  $R_p = 0.7$  fm is the proton radius.

In Fig. 2 we present our results for the total cross section for different values of  $\lambda$  and compare them with experimental data. For references and details see [7].  $\sigma_0$  was assumed to be energy independent [9], important only at lower energies and therefore was not included in our calculations. There is only a small range of values of  $\lambda$  which allow us to describe the experimental data. If, for instance,  $\lambda = 0.4$  the resulting cross section is very flat and clearly below the data, while if  $\lambda = 0.1$  (not shown in figure) the cross section grows very rapidly deviating strongly from the experimental data. The best choice for  $\lambda$  is in the range  $0.25 - 0.30$ , which is exactly the range predicted in theoretical estimates using CGC physics and usually obtained by the saturation models for the  $ep$  HERA data. In [7] we have replaced the IIM dipole cross section by the more modern ones given in [10] but the results do not change very much.




Fig. 2. Energy behavior of the total  $pp/pp\bar{p}$  cross section for different values of  $\lambda$ .

## REFERENCES

- [1] T. K. Gaisser and F. Halzen, Phys. Rev. Lett. **54**, 1754 (1985).
- [2] L. Durand and H. Pi, Phys. Rev. D **40**, 1436 (1989); Nucl. Phys. Proc. Suppl. **12**, 379 (1990); R. M. Godbole, A. Grau, G. Pancheri and Y. N. Srivastava, Phys. Rev. D **72**, 076001 (2005).
- [3] E. Iancu and R. Venugopalan, arXiv:hep-ph/0303204; J. Jalilian-Marian and Y. V. Kovchegov, Prog. Part. Nucl. Phys. **56**, 104 (2006).
- [4] E. Ferreira, E. Iancu, K. Itakura and L. McLerran, Nucl. Phys. A **710**, 373 (2002); A. Kovner and U. A. Wiedemann, Phys. Lett. B **551**, 311 (2003).
- [5] A. D. Martin, R. G. Roberts, W. J. Stirling and R. S. Thorne, Eur. Phys. J. C **23**, 73 (2002).
- [6] J. Bartels, E. Gotsman, E. Levin, M. Lublinsky and U. Maor, Phys. Lett. B **556**, 114 (2003).
- [7] F. Carvalho, F. O. Durães, V. P. Gonçalves and F. S. Navarra, arXiv:0705.1842 [hep-ph].
- [8] E. Iancu, K. Itakura, S. Munier, Phys. Lett. B **590**, 199 (2004).
- [9] See, for example, H. G. Dosch, E. Ferreira and A. Kramer, Phys. Rev. D **50**, 1992 (1994); H. G. Dosch, F. S. Navarra, M. Nielsen and M. Rueter, Phys. Lett. B **466**, 363 (1999).
- [10] V. P. Gonçalves, M. S. Kugeratski, M. V. T. Machado and F. S. Navarra, Phys. Lett. B **643**, 273 (2006); M. S. Kugeratski, V. P. Gonçalves and F. S. Navarra, Eur. Phys. J. C **44**, 577 (2005).